

Aspects of 2-dimensional Elementary Topos Theory

Luca Mesiti

University of Leeds

CT2024 Santiago de Compostela

24/06/2024

1-dimensional elementary topoi

Set is the archetypal elementary topos

$$\begin{array}{ccc}
 A & \xrightarrow{\quad} & 1 \\
 \downarrow \text{subset } \forall i_A & \lrcorner & \downarrow T \\
 X & \overset{\exists! \chi_A}{\dashrightarrow} & \{T, F\} = 2 \\
 & x \mapsto T \text{ iff } x \in A &
 \end{array}$$

$$\forall X \in \mathcal{S}et$$

$$\mathcal{G}_{T,X}: \mathcal{S}et(X, 2) \xrightarrow[\text{pb along } T]{} \mathcal{S}ub(X)$$

is a bijection.

1-dimensional elementary topoi

Set is the archetypal elementary topos

$$\begin{array}{ccc}
 A & \xrightarrow{\quad} & 1 \\
 \downarrow \text{subset } \forall i_A & \lrcorner & \downarrow T \\
 X & \xrightarrow[\exists! \chi_A]{\quad} & \{T, F\} = 2 \\
 & x \mapsto T \text{ iff } x \in A &
 \end{array}$$

$$\forall X \in \mathcal{S}et$$

$$\mathcal{G}_{T,X}: \mathcal{S}et(X, 2) \xrightarrow[\text{pb along } T]{\quad} \text{Sub}(X)$$

is a bijection.

Definition.

Let \mathcal{C} be a 1-category with finite limits. A **subobject classifier** in \mathcal{C} is a map $\tau: 1 \hookrightarrow \Omega$ in \mathcal{C} such that $\forall X \in \mathcal{C}$ $\mathcal{G}_{T,X}: \mathcal{C}(X, \Omega) \rightarrow \text{Sub}(X)$ given by pulling back τ is a bijection.

2-dimensional elementary topoi

In dimension 2, Weber proposed to classify discrete opfibrations.

Their fibres are sets, thus of one dimension higher than the fibres of subsets.

Definition.

A **discrete opfibration** is a functor $p: \mathcal{E} \rightarrow \mathcal{B}$ such that for every $E \in \mathcal{E}$ every $f: p(E) \rightarrow B$ in \mathcal{B} has a unique lifting to E .

$$\begin{array}{ccccc} \mathcal{E} & & E & \overset{\exists! \bar{f}^E}{\dashrightarrow} & f_* E \\ \downarrow p & & \downarrow p & & \downarrow p \\ B & & p(E) & \xrightarrow{f} & B \end{array}$$

Discrete opfibrations in a 2-category are defined by representability.

2-dimensional elementary topoi

Cat is the archetypal elementary 2-topos

Its 2-dimensional classification is given by the **category of elements (Grothendieck construction)**, that exhibits equivalences

$$Cat(\mathcal{B}, Set) \simeq \mathcal{D}OpFib^s(\mathcal{B})$$

between copresheaves and discrete opfibrations with small fibres.

2-dimensional elementary topoi

Cat is the archetypal elementary 2-topos

Its 2-dimensional classification is given by the **category of elements** (**Grothendieck construction**), that exhibits equivalences

$$\mathit{Cat}(\mathcal{B}, \mathit{Set}) \simeq \mathcal{D}OpFib^s(\mathcal{B})$$

between copresheaves and discrete opfibrations with small fibres.

The category of elements is equivalently given by the **comma object**

$$\begin{array}{ccc} \mathcal{E} & \longrightarrow & 1 \\ \downarrow \scriptstyle \begin{array}{l} \forall p \\ \text{disc opfib} \\ \text{small fibres} \end{array} & \swarrow \scriptstyle \text{comma} & \downarrow \scriptstyle 1=\omega \\ \mathcal{B} & \dashrightarrow_{\exists \chi_p} & \mathit{Set} \\ & \text{taking fibres} & \end{array}$$

$$\forall \mathcal{B} \in \mathit{Cat}$$

$$\mathit{Cat}(\mathcal{B}, \mathit{Set}) \xrightarrow[\text{comma along } \omega]{\widehat{\mathcal{G}}_{\omega, \mathcal{B}}} \mathcal{D}OpFib^s(\mathcal{B})$$

is an equivalence of categories.

Definition (M., stronger version of Weber's notion).

Let \mathcal{L} be a 2-category with comma objects, pullbacks along discrete opfibrations and terminal objects. Let P be a fixed pullback stable property for discrete opfibrations. A **good 2-classifier in \mathcal{L}** (w.r.t. P) is a morphism $\omega: 1 \rightarrow \Omega$ in \mathcal{L} such that for every $F \in \mathcal{L}$ the functor

$$\widehat{\mathcal{G}}_{\omega, F}: \mathcal{L}(F, \Omega) \rightarrow \mathcal{D}OpFib(F)$$

given by taking comma objects from ω forms an equivalence of categories when restricting the codomain to the full subcategory $\mathcal{D}OpFib^P(F)$ on the discrete opfibrations that satisfy P .

Dense generators

Definition.

A 2-functor $I: \mathcal{Y} \hookrightarrow_{\text{ff}} \mathcal{L}$ is **dense** if the restricted Yoneda embedding

$$\begin{aligned} \widetilde{I}: \mathcal{L} &\longrightarrow [\mathcal{Y}^{\text{op}}, \text{Cat}] \\ F &\mapsto \mathcal{L}(I(-), F) \end{aligned} \quad \text{is fully faithful.}$$

Equivalently, **every $F \in \mathcal{L}$ is an I -absolute** (i.e. preserved by \widetilde{I})
2-colimit of a diagram that factors through \mathcal{Y} .

Dense generators

Definition.

A 2-functor $I: \mathcal{Y} \hookrightarrow_{\text{ff}} \mathcal{L}$ is **dense** if the restricted Yoneda embedding

$$\begin{aligned} \widetilde{I}: \mathcal{L} &\longrightarrow [\mathcal{Y}^{\text{op}}, \text{Cat}] \\ F &\mapsto \mathcal{L}(I(-), F) \end{aligned} \quad \text{is fully faithful.}$$

Equivalently, **every $F \in \mathcal{L}$ is an I -absolute** (i.e. preserved by \widetilde{I}) **2-colimit of a diagram that factors through \mathcal{Y} .**

Example.

Representables form a dense generator $y: \mathcal{C} \rightarrow [\mathcal{C}^{\text{op}}, \text{Cat}]$ of 2-presheaves. Every 2-presheaf is a weighted 2-colimit of representables.

Reduction of 2-classifiers to dense generators

Theorem (M.).

Let $I: \mathcal{Y} \xrightarrow[\text{ff}]{\hookrightarrow} \mathcal{L}$ be dense and consider $\omega: 1 \rightarrow \Omega$ in \mathcal{L} . **TFAE**:

- (i) ω is a good 2-classifier in \mathcal{L} , i.e. $\forall F \in \mathcal{L} \ \widehat{G}_{\omega, F}$ is an equivalence;
- (ii) $\forall Y \in \mathcal{Y} \ \widehat{G}_{\omega, Y}$ is an equivalence and a certain **operation of normalization** is possible.

$$\begin{array}{ccccc}
 H(C, X) & \longrightarrow & G & & 1 \\
 \psi^{(C, X)} \downarrow & \lrcorner & \downarrow \varphi & & \downarrow \omega \\
 K(C, X) & \xrightarrow{\Lambda_{(C, X)}} & F & \overset{\chi}{\dashrightarrow} & \Omega \\
 & \searrow \widehat{G}_{\omega}^{-1}(\psi^{(C, X)}) & & &
 \end{array}$$



M. 2-classifiers via dense generators and Hofmann-Streicher universe in stacks, [arXiv:2401.16900](https://arxiv.org/abs/2401.16900), 2024.

Problem: colimits in 2-dimensional slices

Theorem (M.).

Let \mathcal{L} be a 2-category and $M \in \mathcal{L}$. Then the 2-functor $\text{dom}: \mathcal{L} /_{\text{lax}} M \rightarrow \mathcal{L}$ is a 2-colim-fibration. As a consequence,

$$\begin{array}{ccc} \text{colim}^W F & & \text{oplax}^{\text{cart}}\text{-colim}^{\Delta^1}(F \circ \mathcal{G}(W)) \\ \downarrow q & = & \downarrow q \\ M & & M \end{array} = \text{oplax}^{\text{cart}}\text{-colim}^{\Delta^1} D^q$$

in the lax slice $\mathcal{L} /_{\text{lax}} M$. Here, D^q is the 2-diagram corresponding to the cartesian-marked oplax cocone associated to q .



M. Colimits in 2-dimensional slices, [arXiv:2305.01494](https://arxiv.org/abs/2305.01494), 2023.

Reduction applied to Cat

Example.

The singleton category **1** is dense in Cat . So we can just look at the classification over **1**, where we clearly have an equivalence.

$$\widehat{G}_{\omega,1}: Cat(1, Set) \xrightarrow{\sim} \mathcal{D}OpFib^s(1) \cong Set$$

We deduce from this trivial observation that the category of elements construction is fully faithful and classifies precisely all discrete opfibrations with small fibres.

$$\begin{array}{ccccc} p^{-1}(B) & \longrightarrow & \mathcal{E} & & 1 \\ \downarrow & \lrcorner & \downarrow p & & \downarrow \omega \\ 1 & \xrightarrow{B} & \mathcal{B} & \xrightarrow{\text{collect fibres}} & Set \\ & \searrow p^{-1}(B) & & & \end{array}$$

Stacks: Grothendieck 2-topoi

Definition (Idea).

A **stack** is a **bicategorical sheaf**. It is a 2-functor $F: \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}at$ such that, for every $C \in \mathcal{C}$ and covering sieve $S \in J(C)$, every assignment

$$\begin{aligned} (D \xrightarrow{f} C) \in S &\longmapsto M_f \in F(D) \\ (D' \xrightarrow{g} D \xrightarrow[f \in S]{f} C) &\longmapsto \varphi^{f,g}: g^* M_f \cong M_{f \circ g} \end{aligned}$$

with the $\varphi^{f,g}$ satisfying the cocycle condition can be **glued into a global** $M \in F(C)$ with coherent isomorphisms $\psi^f: f^* M \cong M_f$.

Stacks: Grothendieck 2-topoi

Definition (Idea).

A **stack** is a **bicategorical sheaf**. It is a 2-functor $F: \mathcal{C}^{\text{op}} \rightarrow \mathcal{Cat}$ such that, for every $C \in \mathcal{C}$ and covering sieve $S \in J(C)$, every assignment

$$\begin{aligned} (D \xrightarrow{f} C) \in S &\longmapsto M_f \in F(D) \\ (D' \xrightarrow{g} D \xrightarrow[f \in S]{f} C) &\longmapsto \varphi^{f,g}: g^* M_f \cong M_{f \circ g} \end{aligned}$$

with the $\varphi^{f,g}$ satisfying the cocycle condition can be **glued into a global** $M \in F(C)$ with coherent isomorphisms $\psi^f: f^* M \cong M_f$.

Moreover F is required to be a sheaf on morphisms, i.e. we can glue matching families of morphisms $f^* X \rightarrow f^* Y$ in $F(D)$ in a unique way into global morphisms $X \rightarrow Y$ in $F(C)$.

A good 2-classifier in 2-presheaves

Let \mathcal{C} be a category and consider $\mathcal{L} = [\mathcal{C}^{\text{op}}, \mathbf{Cat}]$. **Representables form a dense generator** $y: \mathcal{C}^{\text{op}} \rightarrow \mathbf{Cat}$, so we can just look at

$$\widehat{\mathcal{G}}_{\omega, y(\mathcal{C})}: [\mathcal{C}^{\text{op}}, \mathbf{Cat}](y(\mathcal{C}), \Omega) \rightarrow \mathcal{D}OpFib^s(y(\mathcal{C}))$$

We want all these functors to be equivalences of categories. So the **Yoneda lemma forces a good 2-classifier** Ω to be, up to equivalence,

$$\mathcal{C} \xrightarrow{\Omega} \mathcal{D}OpFib^s(y(\mathcal{C}))$$

$\omega: 1 \rightarrow \Omega$ picks the identity on every component.

However, this **Ω is only a pseudofunctor, and it is not clear that it lands in small categories.**

Indexed Grothendieck construction

Theorem (Caviglia–M.).

For every 2-functor $F: \mathcal{A} \rightarrow \mathbf{Cat}$, there is a pseudonatural equivalence

$$\mathbf{OpFib}_{[\mathcal{A}, \mathbf{Cat}]}(F) \simeq \left[\int F, \mathbf{Cat} \right]$$

between split opfibrations in $[\mathcal{A}, \mathbf{Cat}]$ over F and 2-copresheaves on the Grothendieck construction $\int F$ of F .

This restricts to a pseudonatural equivalence

$$\mathbf{DOpFib}^s_{[\mathcal{A}, \mathbf{Cat}]}(F) \simeq \left[\int F, \mathbf{Set} \right].$$

This is a **2-dimensional generalization of the fundamental theorem of elementary topos theory**, in the Grothendieck topos case.



Caviglia and M. Indexed Grothendieck construction, *arXiv:2307.16076*, 2023.

A good 2-classifier in 2-presheaves

Theorem (M.).

$$\begin{aligned}\widetilde{\Omega} : \quad \mathcal{C}^{\text{op}} &\longrightarrow \mathbf{Cat} \\ \mathcal{C} &\mapsto [(\mathcal{C}/\mathcal{C})^{\text{op}}, \mathbf{Set}] \\ (C \xleftarrow{f} D) &\mapsto - \circ (f \circ =)^{\text{op}},\end{aligned}$$

equipped with $\widetilde{\omega}: 1 \rightarrow \widetilde{\Omega}$ that picks the constant at 1 presheaf on every component, **is a good 2-classifier in $[\mathcal{C}^{\text{op}}, \mathbf{Cat}]$** that classifies all discrete opfibrations with small fibres.



M. 2-classifiers via dense generators and Hofmann-Streicher universe in stacks, [arXiv:2401.16900](https://arxiv.org/abs/2401.16900), 2024.

A good 2-classifier in 2-presheaves

The proof actually involves **the bicategorical classification process of $\Omega: \mathcal{C} \mapsto \mathcal{D}OpFib^s(y(\mathcal{C}))$**

$$\widehat{G}_{\omega, y(\mathcal{C})}: \text{Ps}[\mathcal{C}^{\text{op}}, \mathcal{CAT}](y(\mathcal{C}), \Omega) \rightarrow \mathcal{D}OpFib^s_{[\mathcal{C}^{\text{op}}, \mathcal{Cat}]}(y(\mathcal{C})) = \Omega(\mathcal{C})$$

which **is isomorphic to the Yoneda lemma's map** and is thus an equivalence.

A good 2-classifier in 2-presheaves

The proof actually involves **the bicategorical classification process of $\Omega: \mathcal{C} \mapsto \mathcal{D}OpFib^s(y(\mathcal{C}))$**

$$\widehat{G}_{\omega, y(\mathcal{C})}: \text{Ps}[\mathcal{C}^{\text{op}}, \mathcal{CAT}](y(\mathcal{C}), \Omega) \rightarrow \mathcal{D}OpFib^s_{[\mathcal{C}^{\text{op}}, \mathcal{Cat}]}(y(\mathcal{C})) = \Omega(\mathcal{C})$$

which **is isomorphic to the Yoneda lemma's map** and is thus an equivalence.

Idea of the normalization process:

$$\begin{array}{ccc} H^{C,X} & \longrightarrow & G \\ \psi^{C,X} \downarrow & \lrcorner & \downarrow \varphi \\ y(\mathcal{C}) & \xrightarrow{X} & F \end{array}$$

change the fibre $(\psi_D^{C,X})_{D \xrightarrow{f} C}$
into the fibre $(\varphi_D)_{F(f)(X)}$
(fibres of φ are global).

Restricting $\tilde{\Omega}$ to a good 2-classifier in stacks

Consider $i: \mathcal{St}(C, J) \subseteq [\mathcal{C}^{\text{op}}, \mathcal{Cat}]$ the full 2-subcategory on **stacks**.
We restrict $\tilde{\omega}: 1 \rightarrow \tilde{\Omega}$ to a good 2-classifier in stacks.

We proved with a general argument that it is enough to find $\Omega_J \in \mathcal{St}(C, J)$ and $\ell: i(\Omega_J) \xrightarrow{\text{ff}} \tilde{\Omega}$ **chronic** such that, given $\varphi: G \rightarrow F$ a discrete opfibration in $\mathcal{St}(C, J)$ with small fibres

$$\begin{array}{ccc}
 1 & \xrightarrow{\tilde{\omega}} & \tilde{\Omega} \\
 \downarrow \exists i(\omega_J) & & \downarrow \ell \\
 i(\Omega_J) & \xrightarrow{\ell} & \tilde{\Omega}
 \end{array}
 \qquad
 \begin{array}{ccc}
 i(G) & \xrightarrow{\quad} & 1 \\
 i(\varphi) \downarrow & \swarrow \text{comma} & \downarrow \tilde{\omega} \\
 i(F) & \xrightarrow{\quad} & \Omega \\
 \downarrow \exists i(\chi_J) & \nearrow \chi & \downarrow \ell \\
 i(\Omega_J) & & \Omega
 \end{array}$$

Then $\omega_J: 1 \rightarrow \Omega_J$ is a good 2-classifier in $\mathcal{St}(C, J)$.

A good 2-classifier in stacks

Theorem (M.).

$$\Omega_J : \mathcal{C}^{\text{op}} \longrightarrow \text{Cat}$$

$$\begin{aligned} \mathcal{C} &\mapsto \mathcal{S}h(\mathcal{C}/\mathcal{C}, J) \subseteq [(\mathcal{C}/\mathcal{C})^{\text{op}}, \text{Set}] \\ (C \xleftarrow{f} D) &\mapsto - \circ (f \circ =)^{\text{op}}, \end{aligned}$$

equipped with $\omega_J : 1 \rightarrow \Omega_J$ that picks the constant at 1 sheaf on every component, **is a good 2-classifier in $\text{St}(\mathcal{C}, J)$** that classifies all discrete opfibrations with small fibres.

This also solves a problem posed by Hofmann and Streicher when attempting to lift Grothendieck universes to sheaves.



M. 2-classifiers via dense generators and Hofmann-Streicher universe in stacks, [arXiv:2401.16900](https://arxiv.org/abs/2401.16900), 2024.

What's next?

- More examples of 2-classifiers
- A refined notion of elementary 2-topos
- 2-dimensional Grothendieck topologies
- 2-dimensional fundamental theorem of elementary topos theory
- The logic of an elementary 2-topos